

# Global Overview of Ecosystem Services Provided by Riparian Vegetation

TENNA RIIS, MARY KELLY-QUINN, FRANCISCA C. AGUIAR, PARASKEVI MANOLAKI, DANIEL BRUNO, MARÍA D. BEJARANO, NICOLA CLERICI, MARÍA ROSÁRIO FERNANDES, JOSÉ C. FRANCO, NEIL PETTIT, ANA P. PORTELA, OLGA TAMMEORG, PRIIT TAMMEORG, PATRICIA M. RODRÍGUEZ-GONZÁLEZ, AND SIMON DUFOUR

*Fluvial riparian vegetation (RV) links fluvial and terrestrial ecosystems. It is under significant pressure from anthropogenic activities, and therefore, the management and restoration of RV are increasingly important worldwide. RV has been investigated from different perspectives, so knowledge on its structure and function is widely distributed. An important step forward is to convert existing knowledge into an overview easily accessible—for example, for use in decision-making and management. We aim to provide an overview of ecosystem services provided by RV by adopting a structured approach to identify the ecosystem services, describe their characteristics, and rank the importance of each service. We evaluate each service within four main riparian vegetation types adopting a global perspective to derive a broad concept. Subsequently, we introduce a guided framework for use in RV management based on our structured approach. We also identify knowledge gaps and evaluate the opportunities an ecosystem service approach offers to RV management.*


*Keywords:* riparian management, streams, rivers, restoration, riparian ecology, ecosystem functions

**R**iparian vegetation (RV) of fluvial systems is a complex of vegetation units along the river network that is functionally related to the other components of the fluvial system and surrounding area (Naiman et al. 2005). It is a hybrid and open ecotone: It is hybrid because it results from coconstruction by human and natural processes, and it is open because the land alongside fluvial systems interacts with the river and associated processes (Dufour et al. 2019). The riparian zone is therefore characterized by high spatial and temporal variability mainly driven by bioclimatic, geomorphological and land-use conditions, which all change over time under natural and human influences. Riparian vegetation in the context of this article is defined as the vegetation established in the floodplain—that is, the portion of terrestrial landscape from the high-water mark toward the upland, where elevated water tables influence vegetation and soil (Naiman et al. 1993).

Riparian vegetation offers a variety of ecosystem services (ES). The concept of ES has become an important model for linking the functioning of ecosystems to human welfare, which is critical for a wide range of decision-making contexts (Fisher et al. 2009). Equally, there is a plethora of definitions of ES, but the general consensus meaning is the

benefits people obtain from ecosystems or the contributions that ecosystems make to human well-being, after the Millennium Ecosystem Assessment (MEA 2005) and the Common International Classification of Ecosystem Services (CICES) report (Haines-Young and Potschin 2013), respectively. The ES concept, which introduced a new framework for analyzing social–ecological systems, has been advocated as a useful tool that provides a holistic and transparent assessment of impacts on human well-being (e.g., MEA 2005, Fischer et al. 2018), allowing decision-making to take proper account of the value of services from ecosystems (Haines-Young and Potschin 2009). Nevertheless, our ability to draw general conclusions on ES in different ecosystems remains limited (Carpenter et al. 2009).

There is a consensus that there should be a distinction between final ES, which are the outputs of ecosystems (whether natural, seminatural, or highly modified), which directly benefit people and can be measured such as fisheries output per season (provisioning ES) or number of lake visitors per year (cultural ES; Lamothe and Sutherland 2018). Supporting or intermediate ES, have little or no direct benefits to people and involve all biophysical structure and processes that support and maintain ecosystems in a favorable



**Figure 1.** The ecosystems cascade model, which highlights the role of supporting processes and intermediate services in the delivery of final services and the goods and benefits humans derive from riparian vegetation. Source: Adapted with permission from Potschin and Haines-Young (2011).


condition that enables the provision of ES by the ecosystem, such as a lake with good water quality that can support fish that could be the final ES. This distinction is well illustrated in the ES cascade framework of Potschin and Haines-Young (2011) to highlight the position of the CICES classification (figure 1). In the present article, we focus on final ES that most directly affect the well-being of people (figure 1; Haines-Young and Potschin 2013).

Riparian vegetation has the capacity to deliver a disproportionately high amount of ES relative to their extent in the landscape (e.g., Sweeney and Newbold 2014) because of their ecotone characteristics and the ecological functions of RV (Capon et al. 2013). However, riparian vegetation is under significant pressure from a range of anthropogenic activities, such as alteration of disturbance regime, stream-flow regulation by dams, pollution, land-use change, timber harvesting, water diversion, mining, deforestation, and from invasive species (figure 1; e.g., Goodwin et al. 1997, Poff et al. 2011). In Europe, it has been estimated that 80% of natural riparian habitats has disappeared during the past 200 years (Naiman et al. 1993). The loss of riparian vegetation is generally immense in developed countries; for example, it has declined by 85%–95% in California, Arizona, and New Mexico, with most losses attributed to grazing (NRC 2002). Conversely, increasing effort is being undertaken to recover RV with varying success depending on the restoration (e.g.,

hydrogeomorphic, active plant introduction, floodplain conversion, invasive species and grazing control (González et al. 2015).

Riparian zones and their vegetation have been investigated from a range of perspectives covering multiple scientific and applied disciplines such as hydrology, biology, geography, remote sensing, management, and restoration (e.g., González et al. 2015). Therefore, knowledge on structure and function of riparian vegetation is distributed among a wide range of fields and disciplines (see Dufour et al. 2019). Several studies have documented how RV is key for specific ES, but few have attempted to document the full range of ES it provides. An important step forward is therefore to convert the existing knowledge into an overview more easily accessible and directly applicable for decision-making and management of riparian vegetation—a task undertaken in this article.

The general objective of this article is to present an overview of ES provided by RV. More specifically, we adopt a structured approach to identify the range of ES, to describe their characteristics, and to rank the importance of each service. We evaluate each service within four main riparian vegetation types structured by local soil moisture and woodiness within a global perspective to derive a broad concept. The key tasks in this article were, therefore, first, to compile a comprehensive checklist of ES provided by RV



**Figure 2.** Diagram showing the four main riparian vegetation types structured along two main factors: cover of wood and local soil moisture.

and, second, to synthesize the knowledge on these ES from the literature. On the basis of the structured approach, we introduce a guided framework for use in riparian vegetation management. We also seek to identify key knowledge gaps and conclude the article by evaluating the opportunities an ES approach offers to riparian vegetation management and restoration.

### Study approach

In this article, we used the groups of final ES described in Maes and colleagues (2016): provisioning, regulating and maintenance, and cultural ES. Provisioning services are the physical products directly obtained from the RV (e.g., timber, seeds, and harvestable genes), regulating and maintenance services incorporate those that both directly (e.g., pollutant capture, carbon sequestration) and indirectly (e.g., regulation of decomposition, climate, and hydrology) sustain environmental quality. Cultural services include tangible recreational uses (e.g., walking along a river) or less tangible benefits such as aesthetic and spiritual benefits and educational values. In fact, the most recent version of CICES (version 5.1; Haines-Young and Potschin 2018) stresses that all ES have an inherent cultural value, but cultural services should be treated as an independent group, as was also undertaken in this article.

We used the CICES framework (CICES version 5.1; Haines-Young and Potschin 2018) to identify the ES provided by RV. We described the characteristic of each ES and included the underlying processes underpinning the ES delivery. We also described the goods and benefits provided by each ES. The characterization of each ES was derived from relevant scientific literature and complemented with empirical information. The final selection of the most important literature referenced in the article was decided by experts in the author group on each particular ES.

A key task of this article was to rank the importance of each ES provided by RV on the basis of the spatial and temporal extent of each ES—that is, how widespread is the ES provision and how often is it occurring. In order to acknowledge that the importance of each ES may vary substantially depending on the type of RV, we ranked the ES importance within each of four broad groupings of RV. The two criteria used for defining these four RV groups were the woodiness of the dominant vegetation (whether herbs or grass, woody shrubs, or trees) and local soil moisture (wet or dry; figure 2). The importance of woody and nonwoody RV for ES was discussed and summarized by Sweeney and Newbold (2014), with a differential provision of services (e.g., nutrient and sediment dynamics) depending on riparian species woodiness. The importance of local soil moisture in structuring

**Table 1. Definitions of the categories of relative importance of ecosystem services (ES) based on the spatial scale at which an ES works (local to global), and the temporal scale of goods and benefits provided by an ES (uncommon to common).**

Temporal scale	Spatial scale			
	Global	Regional	Local	Unknown
Common	High	High	Medium	Unknown
Less than common	High	Medium	Low	Unknown
Uncommon	Medium	Low	Low	Unknown
Unknown	Unknown	Unknown	Unknown	Unknown

Note: The definitions are based on expert opinion and use of scientific literature. The definitions are used to populate table 2, and the color coding is also used in table 3.

and therefore the functioning of RV is well described and includes effects on plant species richness and diversity, plant dispersal and nutrient dynamics (e.g., Nilsson and Svedmark 2002). The selected soil moisture types in our overview capture a representative gradient of conditions from permanently dry aerated soil to temporarily waterlogged soil, to temporary wetlands with surface water to permanent riparian wetlands (figure 2). The four extreme RV types that result from the combination of the two gradients (woodiness and soil moisture) are therefore herbs or grass, dry forest, wet forest, and riparian wetland (figure 2). We have focused on four points in the two-dimensional space characterized by woodiness and local soil moisture, being aware of all intermediate occurring vegetation communities that may differ in ES provision. In addition, environmental settings such as fluvial geomorphology, hydrologic regime, width of the RV and climatic context all would be important but are not separately included in this framework.

We derived the relative importance for each provisioning, regulating, and maintenance ES within each RV group by using the relevant scientific literature and expert opinion of the authors of this article (table 1). For cultural ES, we did not assign relative importance because of a lack of data to support such assessment. A description of each ES is given below, while a synthesis of the ES characterization and ranking is given in table 2.

### Provisioning services from riparian vegetation

In this section we provide an overview of provisioning services from riparian vegetation.

**Biomass.** Biomass production in riparian areas for fuel for heating and green biogas production can be substantial. For example, in the Pacific Northwest, a range of riparian tree species in a lowland floodplain had a density production of 27,000 stems per hectare (ha) in active floodplains and biomass as high as 540 tons dry weight per ha over a 3-year period (Balian and Naiman 2005). Lower values of 54.4 tons per ha (above- and belowground biomass) have been reported under optimal conditions from a riparian forest in Southeast Asia, dominated by *Populus euphratica* Oliv. (Theversus et al. 2012). Biomass production from shrubby vegetation such as certain willow species can be

equally significant. Walczak and colleagues (2018) referred to figures from the United Kingdom, Poland, and Sweden that range from 8 to 12 tons per ha (fresh mass) and up to 20 tons per ha under favorable conditions. Grassy biomass (residual) from managed riparian areas is also considered a provisioning service. Residual biomass from publicly managed floodplains of the Dutch Rhine tributaries was estimated at 370,953 tons dry mass of biomass, of which 87% was grassy biomass (Bout et al. 2019). According to the authors, this was equivalent to an estimated 353 terajoules of heat from the woody biomass and 15 million cubic meters of green biogas from grassy biomass. Taller grasses such as *Phragmites* or *Arundo* are harvested in some areas for biomass (i.e., paludiculture; e.g., Brix et al. 2014) and their stems have been used for thatching and guide agricultural seedlings in many parts of the world.

Food outputs from riparian zone include herbs, berries (elderberries, blackberries, huckleberries, and saskatoon), and, to a lesser extent, mushrooms. In the present article, again, these are alluded to in the literature but do not appear to be widely used, so their importance is limited at local scale and is therefore assessed to be low (see tables 1 and 2).

**Genetic resources.** Genetic resources in RV include any genetic material, such as seeds and spores that could be harvested, wild plants used for crop breeding and genetic information from plant material used to extract genes. Among the wild crop relatives in riparian vegetation, two climber species, grape vine, and hops are used globally in the production of economically and culturally emblematic wine and beer, respectively. In both cases, wild populations are being used and are increasingly recognized for breeding commercial varieties of these species (Patzak et al. 2010). In fact, *Vitis vinifera* L. ssp. *sylvestris* (Gmelin) Hegi, the European wild grape and ancestor of cultivated grapevine varieties (*V. vinifera* L. ssp. *vinifera*), is the sole wild grapevine species existing in Europe. Wild hops (*Humulus lupulus* L.) in riparian areas are potential new germplasm to expand the variability of genetic resources for hop breeding (Patzak et al. 2010).

As a complementary benefit, the genetic resources of crop-wild relatives also have the potential to improve disease resistance of cultivated species. In the case of wild grapevine, comparative inoculation studies with several grapevine

**Table 2. Provisioning, regulating and cultural ecosystem services (ES) and the main goods and benefits provided by riparian vegetation.**

ES section	ES division	ES category	ES	Main goods and benefits	Herbs or grass	Dry forest	Wet forest	Riparian wetlands
Provisioning	Biomass	Standing crop	Standing crop of woody biomass	Biomass for fuel	Low	Medium	Medium	Low
			Standing crop of non-woody biomass		Low	Low	Low	Medium
		Wild plants and their outputs	Harvestable volume of wild berries or other	Food	Low	Low	Low	Low
	Genetic material	Genetic materials from all biota	Seeds, spores and harvestable genes	Extract genes for breeding, new products resisting disease	Unknown	High	Unknown	Unknown
Regulation and maintenance	Transformation of biochemical or physical inputs	Filtration or storage	Filtering or storage of particles	Reduction in sediment and toxic particles transport in streams	High	Medium	High	High
		Carbon sequestration	Fixation storage	Reduction in carbon dioxide	Medium	High	High	High
		Chemical conditions of freshwaters	Removal of nutrient in runoff	Reduced pollution and damage costs of nutrient runoff	Medium	High	Medium	High
	Regulation of physical, chemical and biological conditions	Stabilization and control of erosion	Erosion control	Reduction of erosion and sediment loads in streams	High	Medium	Medium	High
		Buffering and attenuation of mass flows	Landslide	Protect human lives and infrastructure	Low	High	High	Low
		Hydrological cycles and water flow maintenance and flood protection	Flow regulation - The capacity of vegetation to retain water and release it slowly	Damage mitigation of extreme flows	Medium	Medium	High	High
		Pollination	Pollination	Contribution to yield of crops	High	High	Low	Low
		Seed and propagule dispersal	Seed and propagule dispersal	Maintain biodiversity in the region	Unknown	Unknown	Unknown	Unknown
		Maintaining nursery populations and habitats	Providing habitats	Nursery habitats; sustaining populations (e.g., of iconic species, or threaten species)	High	High	High	High
		Pest control	Providing habitats for native pest control agents	Reduction in pest damage to crop	High	High	High	Unknown
Climate regulation	Evaporative cooling by urban riparian trees	Temperature control in stream and air	Low	High	High	Low		
Fire regulation	The capacity of riparian vegetation to reduce frequency, spread or magnitude of fires	Reduction in fire damage costs	Unknown	Unknown	Unknown	High		
Cultural (Biotic)	Direct in situ and outdoor interactions with living systems, that depend on presence in the environmental setting	Experiential and physical interaction	Ecological quality to support recreational use	Recreation, fitness; destressing or mental health; nature-based recreation; ecotourism and ecoawareness; bushwalking, bird watching, orienteering. Also for rest, relaxation and refreshment.	NA	NA	NA	NA

Downloaded from <https://academic.oup.com/bioscience/advance-article-abstract/doi/10.1093/biosci/biaa041/5824857> by guest on 08 May 2020

Table 2. Continued.

ES section	ES division	ES category	ES	Main goods and benefits	Herbs or grass	Dry forest	Wet forest	Riparian wetlands
		Scientific	Sites of specific scientific interest	Knowledge about the environment and nature	NA	NA	NA	NA
		Educational	Sites used for conservation activities	Skills or knowledge about environmental management	NA	NA	NA	NA
		Heritage	Sites of cultural importance	Tourism, local identity	NA	NA	NA	NA
		Aesthetic	Area of natural beauty	Artistic inspiration	NA	NA	NA	NA
	Indirect: remote, often indoor interactions with living systems, that do not require presence in the environmental setting	Sacred or religious values	Totemic species or settings of religious interest	Mental well-being. Many riparian areas provide strong religious significance for indigenous groups, such as particular riparian trees.	NA	NA	NA	NA
		Symbolic values	Species, habitats or landscapes that can be used as symbols	Social cohesion, cultural icon Conservation of riparian habitats and keystone species	NA	NA	NA	NA
		Entertainment	Artistic productions	Nature films, books, paintings, draws	NA	NA	NA	NA
		Existence	Natural areas designated as wilderness	Mental or moral well-being; valuing wilderness of rivers and riverine areas	NA	NA	NA	NA
		Bequest	Species and ecosystem settings	Moral well-being; promotion of the sustainability of biocultural identity and of the overall social-ecological system	NA	NA	NA	NA

Note: For each service, we evaluated the relative spatial and temporal importance within four main riparian vegetation types (see the text and figure 1 for more explanation). The ecosystem services and their main goods and benefits were derived from CICES (version 5.1; <https://cices.eu>). Abbreviation: NA, not applicable.

pathogens revealed relatively high levels of resistance in some of the *Sylvestris* spp. accessions (Schröder et al. 2015). Similar application has been developed in the production of *Rubus* spp. berries. Wild *Rubus idaeus* germplasm from riparian areas could potentially be used against raspberry cane disease, which is among the most devastating problems for raspberry production (Hall et al. 2009).

The genetic pool of wild populations of riparian trees such as the black poplar *Populus nigra* provides economically relevant outputs for the development of commercial native trees and for advanced, molecular breeding of these species. Wild populations of *P. nigra* are also being studied to obtain bioenergy from lignocellulosic feedstocks that has the potential to develop as a sustainable source of renewable energy (Allwright et al. 2016). Finally, seeds provided by riparian species are extremely important as genetic material for ex situ conservation of the native genetic resources.

Overall, at the European scale the relative importance of this ES is reported as unknown (e.g., Vidal-Abarca Gutierrez and Suarez-Alonso 2013).

### Regulating and maintenance services

In this section we provide an overview of regulating and maintenance services provided by riparian vegetation.

**Filtration of pollutants and chemical conditions of freshwaters.** Riparian filtration services refer to the control of sediments, nutrients, and pollutants inputs to adjacent water (Lowrance et al. 1984). A large body of scientific literature demonstrates the important role of riparian zones in regulating and improving water quality in streams and rivers (e.g., Jordan et al. 1993, Kuusemets et al. 2001) involving both physical and biological mechanisms. Physical processes include filtering and deposition of sediments and sediment-bound pollutants, such as pesticides, by roots and

stems (Naiman et al. 2005). As much as 75% of sediments transported from uplands to streams has been reported to be physically retained by RV (Cooper and Gilliam 1987). As was recently discussed in a review by Feld and colleagues (2018), the key to efficient reduction of surface runoff of soil particles is to have grass strips acting as mechanical filters.

Riparian zones are also effective sinks for dissolved inorganic nitrogen and phosphorus from surrounding agricultural or urban areas (Naiman et al. 1997), therefore providing a high potential in controlling eutrophication of water bodies. One major mechanism for nitrogen removal is denitrification (Cooper and Gilliam 1987), occurring in riparian microsites with anaerobic conditions and decomposing organic substrate. Plant and microbial assimilative uptake also contribute significantly to inorganic nitrogen and phosphorus removal within land–water interface environments. Inorganic phosphorus is removed by soil adsorption and deposition of phosphorus-bounded sediments. The efficiency of inorganic nitrogen and phosphorus removal in riparian zones varies because of a number of hydrogeomorphological, chemical and biological factors (e.g., Groffman et al. 1992, Hefting et al. 2003) but denitrification rates up to 295 kilograms of nitrogen per ha per year have been measured in riparian zones (Lowrance et al. 1984). Sabater and colleagues (2003) reported that 5%–30% of nitrogen was removed by meter of buffer strip but with no differences between climate or vegetation type (trees versus herbaceous). In terms of inorganic phosphorus, values of 70% to 90% removals have been estimated in vegetated riparian buffer strips (Gascuel et al. 2010). Finally, pesticides and other contaminants can also be effectively removed in riparian zones by attachment to vegetation matter, biological assimilation and accumulation (sequestration), or by metabolic degradation processes (Aguiar et al. 2015).

**Carbon sequestration.** Carbon sequestration refers to the capture and long-term storage of carbon that would otherwise be emitted to or remain in the atmosphere. Riparian forests and wetlands are important carbon sinks, with potential for long-term storage (table 2; e.g., Cierjacks et al. 2010, Suftin et al. 2016). Given the importance of soil and plant C sequestration to ameliorate changes in atmospheric chemistry, conserving undisturbed riparian areas could be an effective strategy to enhance climate-change mitigation in rivers (e.g., Lal 2005). Sequestration rates vary along environmental gradients and for different vegetation types (e.g., Suftin et al. 2016). Carbon reservoirs in riparian forests seems to be positively correlated with available moisture and negatively correlated with maximum temperature (Suftin et al. 2016, Dybala et al. 2019), although no clear patterns concerning the effects of climate and geological setting in carbon sequestration have yet emerged from the literature. Research on riparian carbon storage and sequestration is scarce and most studies are from South and North America while European studies are even more limited. The median carbon stock in riparian biomass was estimated to be 63 tons

of carbon per ha with the highest values of 318–487 tons of carbon per ha from mature temperate forests in North and South America, and those values were considered to be comparable to the highest estimates for any forest biome (Dybala et al. 2019). From Europe, Cierjacks and colleagues (2010) reported carbon stocks of 474 tons of carbon per ha for mature riparian woods and 212 tons of carbon per ha for meadows and reeds in Danube floodplains. However, the relative importance of distinct riparian compartments for carbon storage and the variations across scales, vegetation types, geological and climate settings are still unknown.

**Erosion control.** Erosion control refers to the reduction of the weathering away of soil and thereby the inputs of soil particles together with nutrients and carbon to water bodies. The soil-stabilizing effect of the plants is particularly relevant during events of intense rainfall and snowmelt (e.g., Larsen 2017). Species composition, root architecture, and woodiness influences control of erosion and riverbank stability (Simon and Collison 2002, Feld et al. 2018). The most effective erosion control seems to result from mixed stands of riparian woody and nonwoody species (Simon and Collison 2002), but as was recently discussed in a review by Feld and colleagues (2018), grass strips are the key to efficient reduction of surface runoff of soil particles.

The reduction and fragmentation of riparian forests, particularly in mountainous areas, endangers the ES of prevention and control of landslides (Larsen 2017). As an extreme example of the importance of keeping riparian zones forested is the La Purisima storm that hit the Panama at the end of 2010. The lack of bank-stabilizing effects of riparian tree roots due to riparian deforestation has been suggested to be one of the causes for more than 500 landslides during this event (Larsen 2017). Overall, this ES is of medium to high importance (table 2).

**Flow regulation.** Floodplains and riparian areas have long been recognized for delivering significant positive flow regulation services involving reduced frequency and magnitude of flooding, augmented low flows, and reduced stream flow and runoff (table 2; Thomas and Nisbet 2007, Rak et al. 2016). Physically, as floodwater flows through a vegetated area, the plants resist the flow, reduce flow velocity, and dissipate the energy, increasing the time available for water to infiltrate into the soil and be stored, which enhances groundwater recharge and results in a delay and reduction in magnitude of downstream flood peaks and reduced riverbank erosion during heavy precipitation events. Increased hydrologic roughness due to vegetation and tree cover further reduces flood peaks. The potential importance is substantial as the damage cost of flooding or drying of urban or agricultural areas is high. For example, the costs for the flood damage caused by an intense rain event (cloudburst) in Copenhagen on 2 July 2011 amounted to 1 billion euros (Leonardsen 2012) and may have been significantly reduced by properly conserving

vegetated upstream riparian areas. Furthermore, the slow release of water from riparian areas during dry periods is important for the ecological health of streams and downstream recipients, as well as for potential crop irrigation in the surrounding areas (Keesstra et al. 2018).

**Pollination and seed dispersal.** Plant regeneration is essential for maintaining biodiversity and ecosystem functioning in ecosystems but may be threatened by human disturbance. Pollination and seed dispersal are the most threatened processes of plant regeneration because any disturbance such as habitat fragmentation or modification by an invasive plant species is likely to change the patterns of seed movement and recruitment. Riparian vegetation provides important nesting and foraging sites (nectar and pollen) for pollinators, and proximity to such habitats has been found to increase pollinator species richness, crop visitation rates, and pollination success (e.g., Garibaldi et al. 2014, Petersen and Nault 2014). Vegetated riparian areas and wetlands support generally higher richness and diversity of pollinator species than dry adjacent lands, especially those dominated by monoculture (e.g., Ricketts 2004, Munyuli et al. 2013). Riparian vegetation may also play a role in seed dispersal across landscapes. However, the significance of this ES provided specifically by riparian vegetation is largely unknown (table 2).

**Maintenance of nursery populations and habitats.** Riparian zones can host highly valuable natural habitats and act as nursery areas. A nursery is defined as a habitat that contributes more than the average, compared with other habitats, to the production of individuals of a particular species that recruits to adult populations (Beck et al. 2001).

Several species of small and large flagship mammals are specialized to inhabit and reproduce in riparian areas (e.g., otter; Prenda et al. 2001), and these areas represent important foraging areas for insectivorous bats (Grindal et al. 1999). Riparian vegetation also provides habitat for resident and migratory birds (e.g., waders, ducks, and herons; e.g., Gillies and Clair 2008), and many species of reptiles and amphibians. Nearly 70% of vertebrate species in a region will use riparian habitats in some significant way during their life cycle (e.g., Naiman et al. 1993). Undisturbed or well conserved riparian areas also positively affect fish productivity (Tomscha et al. 2017) and the presence and spawning of target fish species with commercial and recreational interest such as salmonids (e.g., Bilby et al. 2003). In addition, RV subsidies as leaf litter are especially important for aquatic food webs and in the absence of autochthonous primary production, can be the major carbon source for aquatic biota (e.g., Pettit et al. 2012). Similarly, woody debris in the stream provides habitat and shelter for aquatic organisms and the exposed roots of riparian trees are the spawning substrate and larval habitat for some stream fish species (e.g., Pettit et al. 2013). Riparian vegetation also sustains benthic and riverine invertebrate richness (Malmqvist 2002), and many semiaquatic organisms, such as salamanders, depend on

both aquatic and terrestrial habitats to complete their life cycle and maintain viable populations (Semlitsch 1998).

**Pest control.** Natural control of plant pests in agroecosystems is provided by predators and parasitoids, such as birds, bats, spiders, ground beetles, lady beetles, lacewings, flies, and wasps, as well as entomopathogenic organisms (e.g., fungi, bacteria, nematodes). Habitat requirements for natural enemies include several ecosystem properties often encompassed by riparian systems: supplementary food resources (e.g., alternate hosts or prey), complementary food resources (e.g., pollen, honeydew, nectar), microclimatic conditions, and overwintering or aestivation shelters and refuges (e.g., Jonsson et al. 2008). Riparian vegetation can provide refuges and other resources to natural enemies, and serve as corridors for their dispersion between other noncrop habitats and into crop fields (e.g., Luke et al. 2018). There are several examples worldwide providing evidence of the relevance of pest-regulating services by riparian vegetation. One study by Maisonneuve and Rioux (2001) found that the proportion of pest species decreased with the complexity of riparian vegetation structure, while insectivorous species increased in abundance in woody riparian strips. Stockan and colleagues (2014) reported the highest density and species diversity of generalist predators (Coleoptera, Carabidae) in unmanaged riparian margins. However, Gray and Lewis (2014) observed that riparian forests 30–50 meters wide adjoining oil palm plantations in Malaysian Borneo were unlikely to provide a pest control service. Clearly, the characteristics of riparian vegetation and associated environmental conditions that influence the pest control service require further study.

**Regulation of microclimate.** Riparian vegetation can exert considerable influence on the local microclimate (Chen et al. 1999) with dense, closed canopies reducing evaporation, reducing wind speed and maintaining high relative humidity. In riparian areas, the stream flow regime and groundwater will result in surface soils with high moisture content. Shading from RV canopies also results in lower air and water temperatures, therefore alleviating the heat stress, which is related with public health (e.g., Kristensen et al. 2014). Riparian vegetation is important for temperature and light control within streams (Capon and Pettit 2018). Trees on the river edge provide shade that can reduce instream primary production and water temperature (e.g., Ryan and Kelly-Quinn 2016, Kristensen et al. 2014), the latter with positive effect on dissolved oxygen. This microclimate regulation is especially evident in dry and semidry areas in which the lushness of riparian trees and shrubs contrasts with the surrounding arid landscape in which vegetation is scarce.

**Fire effects mitigation.** Riparian zones can act as a natural barrier to limit the spread and spatial extent of upland wildfires (Pettit and Naiman 2007). Riparian systems tend to differ from adjacent uplands in moisture regime, topography,

microclimate, soils, and vegetative structure, resulting in higher fuel moisture content, relative humidity, and lower wind speeds. Therefore, fires are generally less frequent and of lower intensity in riparian zones than in upland forests and grasslands but vary according to region and forest types. For instance, in prairie grasslands and deciduous riparian woods fire return intervals are periodic and can range from 10 to 30 years (e.g., Dwire and Kauffman 2003). However, in drylands, especially in small order streams or under dry prefire conditions riparian woods can turn into corridors for fire movement (Petit and Naiman 2007).

With changing climate follows increasing risk in many regions of catastrophic fires, so managing this risk while conserving biodiversity is a major challenge. Dense natural riparian vegetation in most cases creates a buffer zone to the stream, which will limit terrestrial fire spread and protect stream ecosystems from fire effects (Bisson et al. 2003). Riparian zones create refugia for fire-sensitive species in a matrix of more fire-prone uplands. The benefits of riparian areas for fire protection have been recorded in diverse climatic environments such as temperate forest in the United States, where, after a severe fire, tree mortality was 37% in the upland area while no trees were killed in the adjacent riparian zone, and there was no loss of diversity of riparian species (Elliot et al. 1999). In a tropical fire-prone savanna in Central America, fire rarely penetrates far into the adjacent riparian forest, and fire damaged trees are only found on gentler slopes near the savanna–forest boundary (Kellman and Tackaberry 1993). However, fragmentation and invasion by exotic fire-tolerant plants (e.g., *Arundo*) can increase fire risk in riparian areas (Busch and Smith 1995), and poorly managed riparian areas that been invaded by fire tolerant species such as grasses *Acacia* or *Eucalyptus* spp. can create severe fire hazards that will actually promote the spread of wildfires.

**Evaluation of the importance of provisioning, regulating, and maintenance services.** When we compared the importance of the provisioning and regulating ES across the four vegetation types given in table 2, we found that out of the 16 services provided, 12 services had at least one high ranking across the four vegetation types, and 6 had medium or high importance across all vegetation types (table 3; filtration or storage, chemical condition of freshwater, fixation storage, erosion control, flow regulation, and providing habitats). Three other services were mainly associated with two vegetation types that are forest and wet forest providing standing crop of woody biomass and climate regulation, and dry woodland together with herbs or grass providing mainly pollination (table 3). In table 3, we ranked the ES provided by RV such that the highest ranked service is the service with highest importance in most of the four vegetation types. It is clear that presence of any of the four vegetation types in the riparian area will provide several ES, but also that forest and riparian wetland will provide more than herbs or grass.

### Cultural services from riparian vegetation

Cultural services are considered “non-material benefits people obtain from ecosystems through spiritual enrichment, cognitive development, reflection, recreation, and aesthetic experiences” (MEA 2005), which have extended the original cultural and recreation categories of Constanza and colleagues (1997). Despite being increasingly recognized as key to ecosystem conservation and unavoidable in the general valuation of ES, apart from recreation services, this broad category is frequently overlooked because of its intangible and subjective nature and because of a lack of methodological frameworks to quantify their value in monetary means (Kumar 2010, Daniel et al. 2012). The CICES framework stresses that all ES have an inherent cultural value, but these services should be treated as an independent section. CICES classifies cultural services into two broad divisions and respective categories: direct cultural services, which cover outdoor interactions with living systems (experiential and physical use of plants, animals, and landscapes; scientific; educational; cultural heritage; aesthetic) and indirect cultural services, which rely mostly on remotely indoor interactions with the environmental setting (sacred or religious values, symbolic values, entertainment, existence, bequest). These services can involve individual species, communities, habitats, and whole ecosystems. In table 2, the direct and indirect cultural services of RV are listed.

**Direct and indirect cultural services.** Riparian areas and vegetation provide opportunities for researching nature *in situ*. They can function as outdoor laboratories for students or local communities through the development of environmental education and citizen science projects by schools, research centers and local associations that may be complemented by experiences in classrooms or science centers. The time scale in which this cultural service is provided can be quite varied, ranging from a single day (e.g., guided visits or activities) to an entire year (long-term education projects). The benefits of outdoor learning experiences have been documented in environmental education literature, and rivers and riparian areas are not an exception. Studies have reported increased environmental awareness and the integration of knowledge of different subject areas from outdoor studies (e.g., Bouillion and Gomez 2001, Overholt and MacKenzie 2005).

The inspiring aesthetic value of rivers and riparian areas are well documented in a comprehensive body of art works (e.g., paintings and drawings) dating back several hundreds of years and in the numerous tourists that visit, for example, the Camargue, the Danube delta, and the Coto de Doñana to enjoy the beauty of nature. The near natural and most diverse sections of rivers are more attractive to people because of a high sense of wilderness (Brown and Daniel 1991, Bowker and Bergstrom 2017). These areas are often in the upper reaches of rivers and are connected to the cultural services associated with mountains and nearby forests (e.g., recreation or exercise in the form of forest

**Table 3. Ecosystem services provided by riparian vegetation, distributed across four main vegetation types, and ranked from high to low importance following definitions of high, medium, and low given in table 1.**

Ecosystem service	Herbs/grass	Forest	Wet forest	Wetlands	ES Importance High to Low
Providing habitats	High	High	High	High	High to Low
Filtering/storage of particles	High	Medium	High	High	
Fixation storage	Medium	High	High	High	
Erosion control	High	Medium	High	High	
Providing habitats for native pest control agents	High	High	High	Unknown	
Flow regulation - The capacity of vegetation to retain water and release it slowly	Medium	Medium	High	High	
Removal of nutrient in runoff	Medium	High	Medium	High	
Landslide	Low	High	High	Low	
Pollination	High	High	Low	Low	
Evaporative cooling by urban riparian trees	Low	High	High	Low	
Standing crop of woody biomass	Low	Medium	Medium	Low	
Seeds, spores and harvestable genes	Unknown	High	Unknown	Unknown	
The capacity of riparian vegetation to reduce frequency, spread or magnitude of fires	Unknown	Unknown	Unknown	High	
Standing crop of non-woody biomass	Low	Low	Low	Medium	
Harvestable volume of wild berries or other	Low	Low	Low	Low	
Seed and propagule dispersal	Unknown	Unknown	Unknown	Unknown	

walks), and cultural heritage and sense of place (Zandersen and Tol 2009).

Riparian vegetation can also provide a sense of continuity and understanding of our place in the universe, which is expressed through ethical and heritage values (Arts et al. 2018). Many riparian areas provide strong religious significance for indigenous groups, such as waterholes or particular riparian trees (e.g., Nagajara et al. 2014). In few cultures from northwestern Europe, wetlands have a spiritual significance, being places in which ghosts, witches, and dwarfs live (e.g., Hauck et al. 2013).

**Methodological framework to guide management**

We have provided and discussed a list of ES provided by RV and ranked the importance across four main vegetation types. The compilation of the full list of ES in one paper can be used to guide decision-making in riparian management and restoration. The framework including the full list of ES provided by RV allows the identification of synergies and trade-offs between ES across RV types. For example, regarding synergies, it is easy from a list of ES as given in table 2 to identify any synergies obtained by choosing certain vegetation types in a given restoration. Regarding trade-offs it is clear that RV provides some services but it can also provide some disservices. For example, RV can decrease flood risk downstream by lowering the speed of flood wave propagation but it can also deliver logs downstream that can generate a notably risk for bridges. Again by having a list of ES any trade-offs become easier to identify and assess for RV management. Therefore, the ES and benefits obtained from RV can be maximized by directing management and


restoration toward specific target ES or bundles of services when taken trade-offs and disservices are taken into consideration. Disservices were not included in this overview because evidence for riparian ecosystems disservices in general is still limited as also pointed out in a recent review on existing evidence on ecosystem disservices (Blanco et al. 2019). However, disservices should be considered systematically in future work in order to be included in ES conceptual framework.

In order to make the concept useful in local management, we aimed to provide a general framework for adopting the ES approach to riparian area management and restoration. We provide a flow chart outlining the steps required in guiding management and restoration using information on provisioning, regulating, maintaining, and cultural services as targets (figure 3). The first part is based on the ecosystem settings, which is the identification of local riparian vegetation and associated ES, and on the assessment and ranking of the importance of these services following table 1. The second part is the decision-making process in which managers need to decide whether the target is a set of specific ES or whether it is to maximize the range of ES benefits.

**Knowledge gaps and perspectives**

Several knowledge gaps can be identified on the basis of the overview given in this article. First, we need more knowledge on how the four main vegetation types and species traits specifically support different ES (tables 2 and 3). The ranking of the importance of each ES across vegetation types was based on expert opinion supported by the literature but in many cases further studies are needed to validate these

Downloaded from https://academic.oup.com/bioscience/advance-article-abstract/doi/10.1093/biosci/biaa041/5824857 by guest on 08 May 2020



**Figure 3.** Management framework for determining the environmental settings for the riparian vegetation for a particular region (ecosystem setting) by identifying (ID) local vegetation types and the relevant ecosystem services and, finally, ranking the importance of the relevant ecosystem services in relating it to the different vegetation types (following table 3). Based on the ecosystem setting we suggest, subsequent steps for managers to make best decisions aiming either for target services or for maximizing the number of services provided by riparian vegetation. The steps include the identification of synergies and trade-offs between ecosystem services and the economical and noneconomical valuation of the target single or bundle of ecosystem services provided by the riparian vegetation.

rankings. Moreover, we have only considered the main vegetation types, but many intermediate vegetation types are present and might support different combinations of ES. Second, seed and propagule dispersal, gene resources, and fire protection are highly understudied ES provided by RV. Third, a general issue across all ES is the matter of spatial scale. How much area is needed in order to support and optimize each of the ES? This is not only an unexplored issue in RV management but also in many other ecosystems when considering ES provision (e.g., Sutherland et al. 2016). Fourth, cultural services are important but currently it is difficult to quantitatively value the benefits; therefore, they are harder to include in management planning. And last, a better understanding of who is really benefiting from ES provided by riparian vegetation is also needed. Therefore, more research on assessment, description, valuation and integration of cultural services into a decision-making is needed (e.g., Vidal-Abarca Gutierrez and Suarez-Alonso 2013).

## Conclusions

The severe degradation worldwide of freshwater ecosystems has posed a major threat to ES of riparian areas and their vegetation. This negative trend has continued to increase in centuries and most severely since 1950 even though the economic implications are serious (e.g., due flood damages), and in many places, this negative trend might even be intensified because of climate change (e.g., Capon et al. 2013). Therefore, restoration of floodplain and RV would represent an important practice to mitigate the effects of such degradation and in many places; this is already occurring. Nevertheless, we consider that, currently, most water-related restoration projects just aim to improve habitat or water quality but may miss other important ES. In order to maximize the benefits of these restoration investments we suggest adopting an ES-based decision-making approach that should include RV. Therefore, if a broader perspective on ES of RV is included to help guide riparian management, the multifunctionality of freshwater ecosystems can be protected and the provision of ES recovered or improved and the benefits to society enhanced.

In order to progress in this approach, more knowledge conversion is needed. However, as was pointed out by Dufour and colleagues (2019), although there has been a continuous increase in the

number of publications on RV since the 1990s, the integration of that knowledge across disciplines and sociocultural aspects of RV are still very much understudied. In the present article, we have listed and ranked ES provided by RV and allocated the importance of each provisioning and regulating ES within each of four broad RV types. We also included cultural services, although we could not systematically assess their importance because of the highlighted knowledge gaps. Finally, we provided the first steps for a guided management framework for including ES in local restoration planning of RV. In order to move from the knowledge-based approach provided in this article to the policy tools for prioritizing restoration, we need to advance mapping of ES and perform assessments of their economic value. Despite the current limitations on available information, we believe this article is a useful start for knowledge conversion and future implementation of the ES approach in restoration and management of RV.

## Acknowledgments

Funding was provided by COST Action (CA16208), CONVERGES: Knowledge Conversion for Enhancing Management of European Riparian Ecosystems and Services. We also acknowledge the Forest Research Centre, a research unit funded by Fundação para a Ciência e a Tecnologia I. P. Portugal (The Portuguese Foundation for Science and Technology; FCT), for support through grant no. UID/AGR/00239/2019 and RIVEAL (grant no. PTDC/CTA-AMB/29790/2017). FCA and MRF were funded by FCT under Norma Transitória grants no. DL57/2016 and no. DL57/2016/CP1382/CT0019, respectively. APP was funded by FCT under grant no. SFRH/BD/115030/2016. PMR-G was funded by FCT investigator program no. IF/00059/2015. DB was funded by a “Juan de la Cierva” research contract (Spanish MINECO grant no. FJCI-2016-29856) at Instituto Pirenaico de Ecología, Spain.

## Supplemental material

Supplemental data are available at *BIOSCI* online.

## References cited

- Arts K, Rabelo MTO, de Figueiredo DM, Maffey G, Ioris AAR, Girard P. 2018. Online and offline representations of biocultural diversity: A political ecology perspective on nature-based tourism and indigenous communities in the Brazilian Pantanal. *Sustainability* 10: 3643.
- Balian EV, Naiman RJ. 2005. Abundance and production of riparian trees in the lowland floodplain of the Queets River Washington. *Ecosystems* 8: 841–861.
- Beck MW, et al. 2001. The identification conservation and management of estuarine and marine nurseries for fish and invertebrates. *BioScience* 51: 633–641.
- Bilby RE, Beach EW, Fransen BR, Walter JK, Bisson PA. 2003. Transfer of nutrients from spawning salmon to riparian vegetation in western Washington. *Transactions of the American Fisheries Society* 132: 733–745.
- Bisson PA, Rieman BE, Luce C, Hessberg PF, Lee DC, Kershner JL, Reeves GH, Gresswell RE. 2003. Fire and aquatic ecosystems of the western USA: Current knowledge and key questions. *Forest Ecology and Management* 178: 213–229.
- Blanco J, Dendoncker N, Barnaud C, Sirami C. 2019. Ecosystem disservices matter: Towards their systematic integration within ecosystem service research and policy. *Ecosystem Services* 36: 100913.
- Bouillion LM, Gomez LM. 2001. Connecting school and community with science learning: Real world problems and school–community partnerships as contextual scaffolds. *Journal of Research in Science Teaching* 38: 878–898.
- Bout AE, Pfau SF, van der Krabben E, Dankbaar B. 2019. Residual biomass from Dutch riverine areas: From waste to ecosystem service. *Sustainability* 11: 509.
- Bowker JM, Bergstrom JC. 2017. Wild and scenic rivers: An economic perspective. *International Journal of Wilderness* 23: 22–33.
- Brix H, Ye S, Laws EA, Sun D, Li G, Ding X, Pei S. 2014. Large-scale management of common reed *Phragmites australis* for paper production: A case study from the Liaohe Delta China. *Ecological Engineering* 73: 760–769.
- Brown TC, Daniel TC. 1991. Landscape aesthetics of riparian environments: Relationship of flow quantity to scenic quality along a wild and scenic river. *Water Resources Research* 27: 1787–1795.
- Busch DE and Smith SD. 1995. Mechanisms associated with decline of woody species in riparian ecosystems of the southwestern U.S. *Ecological Monographs* 65: 347–370.
- Capon S, Pettit NE. 2018. Turquoise is the new green: Restoring and enhancing riparian function in the Anthropocene. *Ecological Management and Restoration* 19: 44–53.
- Capon SJ, et al. 2013. Riparian ecosystems in the 21st century: Hotspots for climate change adaptation? *Ecosystems* 16: 359–381.
- Carpenter SR, Mooney HA, Agard J, Capistrano D, DeFries RS, Diaz S, Perrings C. 2009. Science for managing ecosystem services: Beyond the Millennium Ecosystem Assessment. *Proceedings of the National Academy of Sciences* 106: 1305–1312.
- Chen J, Saunders S, Crow T, Naiman RJ, Broszofski K, Mroz G, Brookshire B, Franklin JF. 1999. Microclimate in forest ecosystem and landscape ecology. *BioScience* 49: 288–297.
- Cierjacks A, Kleinschmit B, Babinsky M, Kleinschroth F, Markert A, Menzel M, Ziechmann U, Schiller T, Graf M, Lang F. 2010. Carbon stocks of soil and vegetation on Danubian floodplain. *Journal of Plant Nutrition and Soil Science* 173: 644–653.
- Constanza R, et al. 1997. The value of the world's ecosystem services and natural capital. *Nature* 387: 253–260.
- Cooper JR, Gilliam JW. 1987. Phosphorus redistribution from cultivated fields into riparian areas. *Soil Science Society of America Journal* 51: 1600–1604.
- Daniel TC, et al. 2012. Contributions of cultural services to the ecosystem services agenda. *Proceedings of the National Academy of Sciences* 109: 8812–8819.
- Dufour S, Rodríguez-González P M, Laslier M. 2019. Tracing the scientific trajectory of riparian vegetation studies: Main topics approaches and needs in a globally changing world. *Science of the Total Environment* 653: 1168–1185.
- Dwire KA, Kauffman JB. 2003. Fire and riparian ecosystems in landscapes of the western USA. *Forest Ecology and Management* 178: 61–74.
- Dybala KE, Matzek V, Gardali T, Seavy NE. 2019. Carbon sequestration in riparian forests: A global synthesis and meta-analysis. *Global Change Biology* 25: 57–67.
- Elliott KJ, Hendrick RL, Major AE, Vose JM, Swank WT. 1999. Vegetation dynamics after a prescribed fire in the southern Appalachians. *Forest Ecology and Management* 114: 199–213.
- Feld CK, Fernandes MR, Ferreira MT, Hering D, Ormerod SJ, Venohr M, Gutiérrez-Cánovas C. 2018. Evaluating riparian solutions to multiple stressor problems in river ecosystems: A conceptual study. *Water Research* 139: 381–394.
- Fisher B, Turner RK, Morling P. 2009. Defining and classifying ecosystem services for decision making. *Ecological Economics* 68: 643–653.
- Fischer M, et al. 2018. Summary for Policymakers of the Regional Assessment Report on Biodiversity and Ecosystem Services for Europe and Central Asia of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
- Garibaldi LA, et al. 2014. From research to action: Enhancing crop yield through wild pollinators. *Frontiers in Ecology and the Environment* 12: 439–447.
- Gascuel C, Dorioz JM, Krogstad T, Bechmann M. 2010. Create and manage vegetated buffers at field boundaries. COST Action 869. [www.cost869.alterra.nl](http://www.cost869.alterra.nl).
- Gillies CS, Clair CCS. 2008. Riparian corridors enhance movement of a forest specialist bird in fragmented tropical forest. *Proceedings of the National Academy of Sciences* 50: 1974–1977.
- González E, Sher AA, Tabacchi E, Masip A, Poulin M. 2015. Restoration of riparian vegetation: A global review of implementation and evaluation approaches in the international peer-reviewed literature. *Journal of Environmental Management* 158: 85–94.
- Goodwin CN, Hawkins CP, Kershner JL. 1997. Riparian restoration in the western United States: Overview and perspective. *Restoration Ecology* 5: 4–14.
- Gray CL, Lewis OT. 2014. Do riparian forest fragments provide ecosystem services or disservices in surrounding oil palm plantations? *Basic and Applied Ecology* 15: 693–700.

- Grindal SD, Morissette JL, Brigham RM. 1999. Concentration of bat activity in riparian habitats over an elevational gradient. *Canadian Journal of Zoology* 77: 972–977.
- Groffman PM, Gold AJ, Simmons RC. 1992. Nitrate dynamics in riparian forests: Microbial studies. *Journal of Environmental Quality* 21: 666–671.
- Haines-Young R, Potschin M. 2009. Methodologies for Defining and Assessing Ecosystem Services. Joint Nature Conservation Committee. Project code no. C08-0170-0062.
- Haines-Young R, Potschin M. 2013. The Common International Classification of Ecosystem Services Consultation on Version 4 August–December 2012 Report to the European Environment Agency Contract No EEA/IEA/09/003.
- Haines-Young R, Potschin MB. 2018. Common International Classification of Ecosystem Services (CICES) V5 1 and Guidance on the Application of the Revised Structure. European Environment Agency. <https://cices.eu>.
- Hall H, Hummer K, Jamieson A, Jennings S, Weber C. 2009. Raspberry breeding and genetics. Pages 39–353 in Janick J, ed. *Plant Breeding Reviews*. Wiley.
- Hauck J, Görg C, Varjopuro R, Ratamáki O, Jax K. 2013. Benefits and limitations of the ecosystem services concept in environmental policy and decision making: Some stakeholder perspectives. *Environmental Science and Policy* 25: 13–21.
- Hefting M, Bobbink R, Culawé H. 2003. Nitrous oxide emissions and denitrification in chronically nitrate-loaded riparian buffer zones. *Journal of Environmental Quality* 32: 1194–1203.
- Jonsson M, Wratten SD, Landis DA, Gurr GM. 2008. Recent advances in conservation biological control of arthropods by arthropods. *Biological Control* 45: 172–175.
- Jordan TE, Correll DL, Weller DE. 1993. Nutrient interception by a riparian forest receiving inputs from adjacent cropland. *Journal of Environmental Quality* 22: 467–473.
- Keesstra S, Nunes J, Novara A, Finger D, Avelar D, Kalantari Z, Cerdà A. 2018. The superior effect of nature based solutions in land management for enhancing ecosystem services. *Science of the Total Environment* 610: 997–1009.
- Kellman M, Tackaberry R. 1993. Disturbance and tree species coexistence in tropical riparian forest fragments. *Global Ecology and Biogeography Letters* 3: 1–9.
- Kristensen PB, Kristensen EA, Riis T, Baisner AJ, Larsen SE, Verdonshot P, Baattrup-Pedersen A. 2014. The effect of forest on stream temperature. *Inland Waters* 5: 27–38.
- Kumar P, ed. 2010. *The Economics of Ecosystems and Biodiversity: Ecological and Economic Foundations*. Earthscan.
- Kuusemets V, Mander U, Lõhmus K, Ivask M. 2001. Nitrogen and phosphorus variation in shallow groundwater and assimilation in plants in complex riparian buffer zones. *Water Science and Technology* 44: 615–622.
- Lal R. 2005. Forest soils and carbon sequestration. *Forest ecology and management* 220: 242–258.
- Lamothe KA, Sutherland IJ. 2018. Intermediate ecosystem services: The origin and meanings behind an unsettled concept. *International Journal of Biodiversity Science, Ecosystem Services, and Management* 14: 179–187.
- Larsen MC. 2017. Contemporary human uses of tropical forested watersheds and riparian corridors: Ecosystem services and hazard mitigation with examples from Panama Puerto Rico and Venezuela. *Quaternary International* 448: 190–200.
- Leonardsen L. 2012. Financing climate change adaptation: The Copenhagen case. Pages 415–419 in Otto-Zimmermann K, ed. *Resilient Cities 2. Local Sustainability*, vol 2. Springer.
- Lowrance R, Todd RL, Asmussen LE. 1984. Nutrient cycling in an agricultural watershed: I phreatic movement. *Journal of Environmental Quality* 13: 22–27.
- Luke SH, et al. 2018. Riparian buffers in tropical agriculture: Scientific support effectiveness and directions for policy. *Journal of Applied Ecology* 56: 85–92.
- Maes J, et al. 2016. An indicator framework for assessing ecosystem services in support of the EU Biodiversity Strategy to 2020. *Ecosystem Services* 17: 14–23.
- Maisonneuve C, Rioux S. 2001. Importance of riparian habitats for small mammal and herpetofaunal communities in agricultural landscapes of southern Québec. *Agriculture Ecosystems and Environment* 83: 165–175.
- Malmqvist B. 2002. Aquatic invertebrates in riverine landscapes. *Freshwater Biology* 47: 679–694.
- Millennium Ecosystem Assessment. 2005. *Ecosystems and Human Well-being*. Synthesis Island Press.
- Munyuli MBT, Nyeko P, Potts S, Atkinson P, Pomeroy D, Vickery J. 2013. Patterns of bee diversity in mosaic agricultural landscapes of central Uganda: Implication of pollination services conservation for food security. *Journal of Insect Conservation* 17: 79–93.
- Naiman RJ, Décamps R, Henri D. 1997. The ecology of interfaces: Riparian zones. *Annual Review of Ecology and Systematics* 28: 621–658.
- Naiman RJ, Décamps H, Pollock M. 1993. The role of riparian corridors in maintaining regional biodiversity. *Ecological Applications* 3: 209–212.
- Naiman RJ, Décamps H, McClain ME. 2005. *Riparian: Ecology Conservation and Management of Streamside Communities*. Aquatic ecology series. Elsevier Academic Press.
- National Research Council. 2002. *Riparian Areas: Functions and Strategies for Management*. National Academies Press.
- Nilsson C, Svedmark M. 2002. Basic principles and ecological consequences of changing water regimes: Riparian plant communities. *Environmental management* 30: 468–480.
- Overholt E, MacKenzie AH. 2005. Long-term stream monitoring programs in US secondary schools AU. *Journal of Environmental Education* 36: 51–56.
- Patzak J, Nesvadba V, Krofta K, Henychova A, Marzoev AI, Richards K. 2010. Evaluation of genetic variability of wild hops (*Humulus lupulus* L.) in Canada and the Caucasus region by chemical and molecular methods. *Genome* 53: 545–557.
- Petersen JD, Nault BA. 2014. Landscape diversity moderates the effects of bee visitation frequency to flowers on crop production. *Journal of Applied Ecology* 51: 1347–1356.
- Pettit NE, Naiman RJ. 2007. Fire in the riparian zone: Characteristics and ecological consequences. *Ecosystems* 10: 673–687.
- Pettit NE, Davies T, Fellman JB, Grierson PE, Warfe DM, Davies PM. 2012. Leaf litter chemistry decomposition and assimilation by macroinvertebrates in two tropical streams. *Hydrobiologia* 680: 63–77.
- Pettit NE, Warfe DM, Kennard MJ, Pusey BJ, Davies PM, Douglas MM. 2013. Dynamics of in-stream wood and its importance as fish habitat in a large tropical floodplain river. *River Research and Applications* 29: 864–875.
- Poff B, Koestner KA, Neary DG, Henderson V. 2011. Threats to riparian ecosystems in western North America: An analysis of existing literature. *Journal of the American Water Resources Association* 47: 1241–1254.
- Potschin M, Haines-Young R. 2011. Ecosystem services: Exploring a geographical perspective. *Progress in Physical Geography* 35: 575–594.
- Prenda J, López-Nieves P, Bravo R. 2001. Conservation of otter (*Lutra lutra*) in a Mediterranean area: The importance of habitat quality and temporal variation in water availability. *Aquatic Conservation: Marine and Freshwater Ecosystems* 11: 343–355.
- Rak G, Kozelj D, Steinman F. 2016. The impact of floodplain land use on flood wave propagation. *Natural Hazards* 83: 425–443.
- Ricketts TH. 2004. Tropical forest fragments enhance pollinator activity in nearby coffee crops. *Conservation Biology* 18: 1262–1271.
- Ryan D, Kelly-Quinn M. 2016. Riparian vegetation management for water temperature regulation: Implications for the production of macroinvertebrate prey of salmonids. *Fisheries Management and Ecology* 23: 519–530.
- Sabater SA, et al. 2003. Nitrogen removal by riparian buffers along a European climatic gradient: Patterns and factors of variation. *Ecosystems* 6: 20–30.

- Schröder S, Kortekamp A, Heene E, Daumann J, Valea I, Nick P. 2015. Crop wild relatives as genetic resources: The case of the European wild grape. *Canadian Journal of Plant Science* 95: 905–912.
- Semlitsch RD. 1998. Biological delineation of terrestrial buffer zones for pond-breeding salamanders. *Conservation Biology* 12: 1113–1119.
- Simon A, Collison AJ. 2002. Quantifying the mechanical and hydrologic effects of riparian vegetation on streambank stability. *Earth Surface Processes and Landforms* 27: 527–546.
- Stockan JA, Baird J, Langan SJ, Young MR, Iason GR. 2014. Effects of riparian buffer strips on ground beetles (Coleoptera Carabidae) within an agricultural landscape. *Insect Conservation and Diversity* 7: 172–184.
- Sutfin NA, Wohl EE, Dwir KA. 2016. Banking carbon: A review of organic carbon storage and physical factors influencing retention in floodplains and riparian ecosystems. *Earth Surface Processes and Landforms* 41: 38–60.
- Sutherland IJ, Gergel SE, Bennet EM. 2016. Seeing the forest for its multiple ecosystem services: Indicators for cultural services in heterogeneous forests. *Ecological Indicators* 71: 123–133.
- Sweeney BW, Newbold JD. 2014. Streamside forest buffer width needed to protect stream water quality habitat and organisms: A literature review. *Journal of the American Water Resources Association* 50: 560–584.
- Theversus N, Buras A, Zerbe S, Kühnel E, Abdusalih N, Ovezberdiyeva A. 2012. Structure and wood biomass of near-natural floodplain forests along the Central Asian rivers Tarim and Amu Darya. *Forestry* 85: 193–202.
- Thomas H, Nisbet TR. 2007. An assessment of the impact of floodplain woodland on flood flows. *Water Environ Journal* 21: 114–126.
- Tomscha SA, Gergel SE, Tomlinson MJ. 2017. The spatial organization of ecosystem services in river-floodplains. *Ecosphere* 8: e01728.
- Vidal-Abarca Gutierrez MR, Suarez-Alonso ML. 2013. Which are what is their status and what can we expect from ecosystem services provided by Spanish rivers and riparian areas? *Biodiversity Conservation* 22: 2469–2503.
- Walczak N, Walczak Z, Kaluza T. 2018. The Impact of shrubby floodplain vegetation Growth on the discharge capacity of river valleys. *Water* 10: 556.
- Zandersen M, Tol RS. 2009. A meta-analysis of forest recreation values in Europe. *Journal of Forest Economics* 15: 109–130.
- Aguiar TR Jr, Bortolozzo FR, Hansel FA, Rasera K, Ferreira MT. 2015. Riparian buffer zones as pesticide filters of no-till crops. *Environmental Science and Pollution Research* 22: 10618–10626.

---

*Tenna Riis (tenna.riis@bios.au.dk) and Paraskevi Manolaki are affiliated with the Department of Bioscience at Aarhus University, in Aarhus, Denmark. Mary Kelly-Quinn is affiliated with the School of Biology and Environmental Science at University College Dublin, in Dublin, Ireland. Francisca C. Aguiar, Maria Rosário Fernandes, José C., Franco, and Patricia M. Rodríguez-González are affiliated with the Forest Research Centre, in the School of Agriculture at the University of Lisbon, in Lisbon, Portugal. Paraskevi Manolaki is affiliated with the Aarhus Institute of Advanced Studies, at Aarhus University, in Aarhus, Denmark. Daniel Bruno is affiliated with the Instituto Pirenaico de Ecología, in Zaragoza, Spain. María D. Bejarano is affiliated with the Universidad Politécnica de Madrid, in Madrid, Spain. Nicola Clerici is affiliated with the Department of Biology, in the Faculty of Natural Sciences and Mathematics, at the Universidad del Rosario, in Bogotá, Colombia. Neil Pettit is affiliated with the Ecosystems Research Group, in the School of Biological Sciences at The University of Western Australia, in Perth, Australia. Ana P. Portela is affiliated with the Research Center on Biodiversity and Genetic Resources, in the Faculty of Sciences of the University of Porto, in Porto, Portugal. Olga Tammeorg is affiliated with the Estonian University of Life Sciences, in Tartu, Estonia. Priit Tammeorg is affiliated with the University of Helsinki, in Helsinki, Finland. Simon Dufour is affiliated with the University of Rennes, in Rennes, France.*